

Partition Relations for Linear Orders without the Axiom of Choice

03E02, 03E60, 05C63

Thilo Weinert

Department of Mathematics,
Ben-Gurion-University of the Negev,
Israël

Joint work with Philipp Lücke and Philipp Schlicht

3rd SetTop Conference, Fruška Gora,
Tuesday, 21st of June 2016, 12:30-12:55

Folklore ([933Si])

Assume the Axiom of Choice. Then $L \not\rightarrow (\omega^, \omega)^2$ for any linear order L .*

Theorem ([965Kr, Theorem 8] and [971E, Theorem 5])

Assume the Axiom of Choice. Then

$L \not\rightarrow (4, \omega^* + \omega)^3$ and

$L \not\rightarrow (4, \omega + \omega^*)^3$ for any linear orders L .

Theorem ([971E, Theorem 5])

Assume the Axiom of Choice. Then $L \not\rightarrow (5, \omega^ + \omega \vee \omega + \omega^*)^3$ for all linear orders L .*

Question

Assume the Axiom of Choice. Is there a linear order L with $L \rightarrow (4, \omega + \omega^ \vee \omega^* + \omega)^3$?*

Theorem ([976Pr])

The axiom of determinacy of games of reals $\text{AD}_{\mathbb{R}}$ implies that $\omega \rightarrow (\omega)_2^\omega$.

Theorem ([977Ma, 5.1 Metatheorem])

It is consistent from an inaccessible cardinal that $\omega \rightarrow (\omega)_2^\omega$.

Theorem (Donald Martin, [003Ka, Theorem 18.12],
[004JM, 990Ja, 981K])

The axiom of determinacy AD implies that $\omega_1 \rightarrow (\omega_1)_2^{\omega_1}$.

Observation

$\langle {}^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^*, \omega)^3$ for all ordinals α .

Theorem ([981B1])

For every *continuous* colouring χ with $\text{dom}(\chi) = [{}^\omega 2]^n$ there is a perfect $P \subset {}^\omega 2$ on which the value of χ at an n -tuple is decided by its splitting type.

Folklore ([967My, 978Ta])

Every relation on the reals with the property of Baire is continuous on a perfect set.

Observation

$\langle {}^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^*, \omega)^3$ for all ordinals α .

Theorem ([981B1])

For every *Baire* colouring χ with $\text{dom}(\chi) = [^\omega 2]^n$ there is a perfect $P \subset {}^\omega 2$ on which the value of χ at an n -tuple is decided by its splitting type.

Folklore ([967My, 978Ta])

Every relation on the reals with the property of Baire is continuous on a perfect set.

Theorem

Suppose that all sets of reals have the property of Baire.

Then $\langle {}^{\omega}2, <_{lex} \rangle \rightarrow (\langle {}^{\omega}2, <_{lex} \rangle)_n^2$ for all n .

Theorem

Suppose that all sets of reals have the property of Baire.

Then $\langle {}^{\omega}2, <_{lex} \rangle \rightarrow (\langle {}^{\omega}2, <_{lex} \rangle, 1 + \omega^ \vee \omega + 1)^3$.*

Summary

Assume that all sets of reals have the property of Baire. Then

$$\langle {}^{\omega}2, <_{lex} \rangle \rightarrow (\omega + 1)_2^4,$$

$$\langle {}^{\omega}2, <_{lex} \rangle \rightarrow (5, 1 + \omega^* + \omega + 1 \vee \omega + 1 + \omega^*)^4,$$


$$\langle {}^{\omega}2, <_{lex} \rangle \rightarrow (6, 1 + \omega^* + \omega + 1 \vee m + \omega^* \vee \omega + n)^4.$$

combs

candelabra

bouquets

sinistral



dextral



Figure: Combs, Candelabra and Bouquets

- \vec{p} is a *cactus* if and only if s_p divides \vec{p} in a comb of the same chirality and a branch.
- \vec{p} is a *grape* if and only if s_p divides \vec{p} in a comb of the opposite chirality and a branch.
- \vec{p} is an *olivillo* if and only if s_p divides \vec{p} in a bouquet of the same chirality and a branch.
- \vec{p} is a *rose* if and only if s_p divides \vec{p} in a bouquet of the opposite chirality and a branch.
- \vec{p} is a *mistletoe* if and only if s_p divides \vec{p} in a candelabrum and a branch.
- \vec{p} is a *lilac* if and only if s_p divides \vec{p} in a triple of the same chirality and a pair.
- \vec{p} is a *guinea flower* if and only if s_p divides \vec{p} in a triple of the opposite chirality and a pair.




Figure: Seven Pentapetalae, cf. [991HP, 009B&, 010M&, 015St]

Theorem (Lücke & Schlicht)

It is consistent that $\langle^{\omega_1} 2, <_{lex} \rangle \rightarrow (\langle^{\omega_1} 2, <_{lex} \rangle)^2$.

Theorem

Let κ be an infinite initial ordinal and $\alpha < \kappa^+$. Then $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (2 + \kappa^ \vee \omega, \omega^* \vee \kappa + 2)^m$ for all $m \geq 3$.*


Summary

If α is an ordinal, then the following statements hold.

$$\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (5, \omega^* + \omega)^4,$$

$$\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (5, \omega + \omega^*)^4,$$

$$\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (7, \omega^* + \omega \vee \omega + \omega^*)^4.$$

Figure: The functions $\Delta, \delta, \ell, h, \gamma_h$ and β_h


Lemma

For all ordinals α every sextuple within $\langle {}^\alpha 2, <_{lex} \rangle$ contains a cactus, lilac, sinistral bouquet, dextral olivillo or dextral grape (and, by symmetry, a cactus, lilac, dextral bouquet, sinistral olivillo or sinistral grape).


Lemma

Suppose that α is an infinite ordinal and $h : \alpha \hookrightarrow \# \alpha$ is an injection. For every $X \in [{}^\alpha 2]^{\omega^*\omega}$, at least one of the following conditions hold.


- ① There is a candelabrum $\vec{x} = \{x_0, x_1, x_2, x_3\}_{<_{lex}} \in [X]^4$ with $\beta_h(x_1, x_2) < \min(\beta_h(x_0, x_1), \beta_h(x_2, x_3))$.
- ② There is a sinistral comb $\vec{x} = \{x_0, x_1, x_2, x_4\}_{<_{lex}} \in [X]^4$ with $\beta_h(x_1, x_2) < \beta_h(x_0, x_1) < \beta_h(x_2, x_3)$.




$$b(\delta) < b(\varepsilon) < b(\gamma)$$




$$b(\varepsilon) < b(\gamma) < b(\delta)$$




$$b(\varepsilon) \notin b(\gamma) \setminus b(\delta)$$







$$b(\gamma) \notin b(\delta) \setminus b(\varepsilon)$$



$$b(\gamma) < b(\delta) < b(\varepsilon)$$



$$b(\gamma) < \min(b(\delta), b(\varepsilon)) \quad b(\gamma) > \min(b(\delta), b(\varepsilon))$$



$$b(\delta) \notin b(\varepsilon) \setminus b(\gamma)$$

$$b(\gamma) < \min(b(\delta), b(\varepsilon)) \quad b(\gamma) > \min(b(\delta), b(\varepsilon))$$

Theorem

If κ is an infinite initial ordinal and $\alpha < \kappa^+$, then

- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (2 + \kappa^* \vee \kappa + 2 \vee \eta, 5)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee (\kappa 2)^* \vee \kappa 2 \vee \kappa + 2 + \kappa^*, 5)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \kappa + \omega \vee \omega^* + \kappa^*, 6)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega + \omega^* \vee 2 + \kappa^* \vee \kappa + 2, 6)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \kappa \vee 2 + \kappa^* \vee \kappa 2 \vee \omega\omega^*, 6)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee 2 + \kappa^* \vee \kappa + \omega, 7)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \kappa \vee \kappa + 2 \vee 2 + \kappa^* \vee \eta, 7)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \omega + \omega^* \vee (\kappa 2)^* \vee \kappa 2, 8)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \omega \vee \omega^* + \kappa \vee 2 + \kappa^* \vee \kappa + 2 \vee \omega\omega^* \vee \omega^*\omega, 8)^4,$
- $\langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \omega + \omega^* \vee \kappa + 2 \vee 2 + \kappa^*, 9)^4.$

Theorem

If κ is an infinite initial ordinal and $\alpha < \kappa^+$, then

$$\begin{aligned}
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (2 + \kappa^* \vee \kappa + 2 \vee \eta & , 5)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee (\kappa 2)^* \vee \kappa 2 \vee \kappa + 2 + \kappa^* & , 5)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \kappa + \omega \vee \omega^* + \kappa^* & , 6)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega + \omega^* \vee 2 + \kappa^* \vee \kappa + 2 & , 6)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \kappa \vee (\kappa 2)^* \vee \kappa + 2 \vee \omega^* \omega & , 6)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \omega^* + \kappa^* \vee \kappa + 2 & , 7)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \kappa \vee \kappa + 2 \vee 2 + \kappa^* \vee \eta & , 7)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \omega + \omega^* \vee (\kappa 2)^* \vee \kappa 2 & , 8)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\kappa^* + \omega \vee \omega^* + \kappa \vee 2 + \kappa^* \vee \kappa + 2 \vee \omega \omega^* \vee \omega^* \omega & , 8)^4, \\
 \langle^\alpha 2, <_{lex} \rangle \not\rightarrow (\omega^* + \omega \vee \omega + \omega^* \vee \kappa + 2 \vee 2 + \kappa^* & , 9)^4.
 \end{aligned}$$

Question

Which partition relations of the form

$$\langle {}^{\kappa}2, <_{lex} \rangle \rightarrow \left(\bigvee_{\nu < \lambda} K_\nu, \bigvee_{\nu < \mu} L_\nu \right)^n$$

for $n \geq 3$ are (jointly) consistent with ZF (+ DC $_{\kappa}$), and which of the relations for $\kappa = \omega_1$ are provable in the theories ZF + AD + [V = L(\mathbb{R})] and ZF + DC + AD $_{\mathbb{R}}$?

Conjecture (W.)

Assume the Axiom of Determinacy. Then $\langle^{\omega_1}2, <_{lex}\rangle \rightarrow (6, \omega + \omega^* \vee \omega^* + \omega)^4$.

Thank you!



Richard Peter Stanley.

Catalan Numbers.

Cambridge Univ. Press, Cambridge, 2015.



Michael J Moore, Pamela S Soltis,

Charles D Bell, J Gordon Burleigh and

Douglas E Soltis.

Phylogenetic analysis of 83 plastid genes further resolves the early diversification of eudicots.

Proceedings of the National Academy of Sciences, 107(10):4623–4628, 2010.



Birgitta Bremer, Käre Bremer, Mark

Chase, Mike Fay, James Reveal, Douglas E Soltis, Pamela S Soltis and Peter Stevens. An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: Apg iii.

Botanical Journal of the Linnean Society, 2009.



Stephen Craig Jackson and Russell May.

The strong partition relation on ω_1 revisited.

MLQ Math. Log. Q., 50(1):33–40, 2004, doi:10.1002/malq.200310073, http://dx.doi.org/10.1002/malq.200310073.



Akihiro Kanamori.

The higher infinite.

Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003.

Large cardinals in set theory from their beginnings.

-
- Peter John Hilton and Jean Pedersen.
- Catalan numbers, their generalization, and their uses.
- Math. Intelligencer*, 13(2):64–75, 1991, doi:10.1007/BF03024089, http://dx.doi.org/10.1007/BF03024089.
-
- Stephen Craig Jackson.
- A new proof of the strong partition relation on ω_1 .
- Trans. Amer. Math. Soc.*, 320(2):737–745, 1990, doi:10.2307/2001700, http://dx.doi.org/10.2307/2001700.
-
- Andreas Raphael Blass.
- A partition theorem for perfect sets.
- Proc. Amer. Math. Soc.*, 82(2):271–277, 1981, doi:10.2307/2043323.
-
- Alexander Sotirios Kechris, Eugene Meyer
- Kleinberg, Yiannis Nicholas Moschovakis and William Hugh Woodin.
- The axiom of determinacy, strong partition properties and nonsingular measures.
- In *Cabal Seminar 77–79 (Proc. Caltech-UCLA Logic Sem., 1977–79)*, volume 839 of *Lecture Notes in Math.*, pages 75–99. Springer, Berlin, 1981, doi:10.1007/BFb0090236.
-
- Alan D. Taylor.
- Partitions of pairs of reals.*
- Fund. Math.*, 99(1):51–59, 1978, http://matwbn.icm.edu.pl/tresc.php?wyd=1&tom=99.
-
- Adrian Richard David Mathias.
- Happy families.
- Ann. Math. Logic*, 12(1):59–111, 1977.
-
- Karel Libor Prikry.
- Determinateness and partitions.
- Proc. Amer. Math. Soc.*, 54:303–306, 1976, http://www.jstor.org/stable/2040805.
-
- Paul Erdős, Eric Charles Milner and Richard Rado.
- Partition relations for η_α -sets.
- J. London Math. Soc.* (2), 3:193–204, 1971, http://www.renyi.hu/~p_erdos/1971-16.pdf.
-
- Jan Mycielski.
- Algebraic independence and measure.
- Fund. Math.*, 61:165–169, 1967.
-
- Arthur H. Kruse.
- A note on the partition calculus of P. Erdős and R. Rado.
- J. London Math. Soc.*, 40:137–148, 1965, http://jlms.oxfordjournals.org/content/s1-40/1/137.full.pdf.
-
- Waclaw Franciszek Sierpiński.
- Sur un problème de la théorie des relations.
- Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (2), 2(3):285–287, 1933, http://www.numdam.org/item?id=ASNSP_1933_2_2_3_285_0.
-

Choicelessly Partitioning Quadruples

Coda

Yes, we scan!

